

3577 U.S. PTO
03/24/00

Please type a plus (+) sign in inside this box →

EXPRESS MAILING LABEL NO.
EL108178214US

| | | | |
|---|--|--|----------------|
| UTILITY PATENT APPLICATION TRANSMITTAL (Only for new non-provisional applications under 37 CFR 1.53(b)) | | Attorney Docket No. | DM6993 |
| | | First Named Inventor or Application Identifier | |
| | | LAZEWATSKY | |
| | | Express Mail Label No. | EL108178214US |
| | | Express Mailing Date | March 24, 2000 |

| | | | |
|--|--|--|--|
| APPLICATION ELEMENTS See MPEP chapter 600 concerning utility patent applications contents | | ADDRESS TO: Assistant Commissioner for Patents Box Patent Application Washington, DC 20231 | |
| <p>1. <input checked="" type="checkbox"/> Fee (Authority to charge deposit account below.) (Submit an original, and duplicate for fee processing)</p> <p>2. <input checked="" type="checkbox"/> Specification [Total Pages 51] (preferred arrangement set forth below)</p> <ul style="list-style-type: none"> - Descriptive title of the invention - Cross Reference to Related Application (if needed) - Statement Regarding Fed sponsored R&D (if needed) - Reference to Microfiche Appendix (if needed) - Background of the Invention - Brief Summary of the Invention - Brief Description of the Drawings (if filed) - Detailed Description - Claims [No. of Claims 19] - Abstract of the Disclosure | | <p>6. <input type="checkbox"/> Microfiche Computer Program (Appendix)</p> <p>7. Nucleotide and/or Amino Acid Sequence Submissions (if applicable, all necessary)</p> <p>a. <input type="checkbox"/> Computer Readable Copy</p> <p>b. <input type="checkbox"/> Paper Copy (identical to computer copy)</p> <p>c. <input type="checkbox"/> Statement verifying identity of above copies</p> | |
| <p>3. <input checked="" type="checkbox"/> Drawing(s) (35 USC 113 [Total Pages 1])</p> <p>4. <input checked="" type="checkbox"/> Oath or Declaration [Total Pages 1]</p> <p>a. <input type="checkbox"/> Unexecuted (original or copy)</p> <p>b. <input type="checkbox"/> Copy from a prior application (37 CFR 1.63(d) (for continuation/divisional with Box 14 completed)</p> <p>i. <input type="checkbox"/> DELETION OF INVENTORS Signed Statement below at 15 deleting inventor(s) named in the prior application see 37 CFR 1.63(d)(2) and 1.33(b).</p> | | <p>8. <input checked="" type="checkbox"/> Power of Attorney</p> <p>9. <input type="checkbox"/> Information Disclosure Statement (IDS) Cover Letter plus PTO-1449 Copies of IDS Citations</p> <p>10. <input checked="" type="checkbox"/> Preliminary Amendment</p> <p>11. <input checked="" type="checkbox"/> Return Receipt Postcard (MPEP 503) (Should be specifically itemized)</p> <p>12. <input type="checkbox"/> Certified Copy priority Document(s) (if foreign priority is claimed)</p> <p>13. <input type="checkbox"/> Other</p> | |
| <p>- If a CONTINUING APPLICATION, check appropriate box and supply the requisite information:</p> <p><input type="checkbox"/> Continuation <input type="checkbox"/> Divisional <input type="checkbox"/> Continuation-in-part (CIP) <input type="checkbox"/> prior Application No.:</p> | | | |
| <p>15. <input type="checkbox"/> DELETION OF INVENTOR(S) STATEMENT: This application is being filed by less than all the inventors named in the prior application. In accordance with 37 CFR 1.63(d)(2) and 1.33(b), the Assistant Commissioner is requested to delete the number(s) of the following person or persons who are not inventors of the invention being claimed in this application:</p> | | | |
| <p>16. <input checked="" type="checkbox"/> Amend the specification by inserting before the first line the sentence: This application claims the benefit of U.S. Provisional Application No. 60/126,359 filed on March 26, 1999</p> | | | |
| <p>17. <input type="checkbox"/> Cancel in this application original claims ____ of the prior application before calculating the filing. (At least one Original independent claim must be retained for filing purposes.)</p> | | | |
| <p>18. <input type="checkbox"/> Priority of foreign Application No. _____ filed on _____ in _____ (country) is claimed under 35 U.S.C. 119.</p> | | | |

03/24/00
60126359
U.S. PTO

| CLAIMS | (1) FOR | (2) NUMBER FIELD | (3) NUMBER EXTRA | (4) RATE | (5) CALCULATIONS |
|--------|--|------------------|------------------|--------------------------------------|------------------|
| | TOTAL CLAIMS (37 CFR 1.16(c)) | 19- (20) = | 0 | x \$ (18) = | \$0 |
| | INDEPENDENT CLAIMS (37 CFR 1.16(b)) | 4 - (3) = | 1 | 1x \$ (78) = | \$78 |
| | MULTIPLE DEPENDENT CLAIM(S) (if applicable) | | | = \$270 = | |
| | | | | BASIC FEE (37 CFR 1.16(a)) | \$690.00 |
| | | | | TOTAL = | \$768.00 |

19. The Commissioner is hereby authorized to credit overpayments or charge the following fees to Deposit Account No. 04-1928:

a. ☒ Fees required under 37 CFR 1.16.

b. ☐ Fees required under 37 CFR 1.17

20. ☐ Other:

| 21. CORRESPONDENCE ADDRESS | | | | | |
|----------------------------|---|-----------|--------------|----------|--------------|
| NAME | Maureen P. O'Brien | | | | |
| ADDRESS | DuPont Pharmaceuticals Company c/o E. I. du Pont de Nemours and Company Legal - Patents 1007 Market Street | | | | |
| CITY | Wilmington | STATE | Delaware | ZIP CODE | 19898 |
| COUNTRY | U.S.A. | TELEPHONE | 302-992-4528 | FAX | 302-992-4878 |

| 22. SIGNATURE OF ATTORNEY OR AGENT REQUIRED | | |
|---|---------------------------|------------------|
| NAME | Maureen P. O'Brien | REG. NO.: 42,043 |
| SIGNATURE | <i>Maureen P. O'Brien</i> | |
| DATE | <i>March 24, 2000</i> | |

Method for Localization of Blood Clots

FIELD OF THE INVENTION

The present invention relates to a medical diagnostic method and, in particular, to an *in vivo* diagnostic method for detecting a blood clot, such as a pulmonary embolism or a thrombus, employing a radiopharmaceutical contrast agent and volume rendering of single photon emission computed tomography (SPECT) images.

BACKGROUND OF THE INVENTION

Pulmonary embolism is a condition of the lung that emerges when a portion of a blood clot (i.e., thrombus) growing pathologically within a patient breaks off (i.e., embolizes) and travels to the lung. In many instances, the condition itself is immediately life-threatening. However, even when the condition is not immediately life-threatening, a patient presenting with symptoms characteristic of pulmonary embolism must be properly diagnosed to assure that the symptoms do not represent other diseases. Accordingly, detection and localization of pulmonary embolism are critical to insure that the patient receives the appropriate care.

Previously, a technique for diagnosing tumors has been developed. The technique involves localizing a contrast agent at the tumor and obtaining a series of image slices of the tumor using single photon emission computed tomography (SPECT). The image slices are then individually inspected by a physician. As a result, the process is time consuming and expensive.

To improve the ability to diagnose tumors using SPECT, a volume rendering technique has been developed for displaying SPECT data derived from a complete set of image slices through the tumor. According to this technique, a three-dimensional matrix of data is assembled from the image slices. The three-dimensional matrix of data is then scanned along an array of parallel lines at a given angle with respect to the tumor.

For each parallel line, the value of the most intense pixel along the parallel line is determined and assigned to a pixel in a two-dimensional array whose position corresponds to the position of the corresponding parallel line in the array of parallel lines. The process is repeated for a series of angles over 360° to produce a series of two-dimensional images. When the series of two-dimensional images are displayed sequentially, a rotating view of the most intense pixels is produced.

In spite of the foregoing, the utility of SPECT as a tool for diagnosing pulmonary embolism remained limited. The limited use of SPECT in connection with pulmonary embolism is due, at least in part, to the fact that the normal anatomy of the thorax is complex. As a result, structures highlighted by the contrast agent are variable, often in a pattern that is unfamiliar to physicians, and without the normal identifying landmarks. Thus, the location of the thrombus and the extent of disease was expected to be difficult to ascertain from the SPECT images, even if volume rendering techniques were employed.

Accordingly, it would be highly beneficial to provide a method for identification and localization of pulmonary embolism using SPECT wherein a three-dimensional representation of a thrombus is obtained. The three-dimensional representation of the thrombus should enable a physician to more clearly, accurately, and efficiently determine the extent of disease. Accordingly, the present invention should provide a significant qualitative improvement in the ability of a naive physician to identify and localize a thrombus.

SUMMARY OF THE INVENTION

The shortcomings associated with the known methods for localization of blood clots are overcome to a large degree by a method in accordance with the present invention. The method according to the present invention comprises the step of localizing a radiolabelled compound at a thrombus by

administering a radiopharmaceutical compound to the patient. Two-dimensional images representing a physical property associated with the radiolabelled thrombus, such as single photon emission computed tomography (SPECT) images, are then acquired and assembled into a three-dimensional matrix of data. The three-dimensional matrix of data is then scanned along an array of parallel lines to determine a maximum value along each line. The maximum value along each line is then assigned to a pixel in a two-dimensional array, where the relative position of the pixel in the two-dimensional array corresponds to the relative position of the line in the array of parallel lines. The three-dimensional matrix of data is optionally scanned along additional arrays of parallel lines to produce a series of images of the thrombus as viewed from different angles. The series of images can be displayed sequentially to produce a rotating view of the thrombus.

BRIEF DESCRIPTION OF THE DRAWINGS

The foregoing summary, as well as the following detailed description of the preferred embodiments of the present invention, will be better understood when read in conjunction with the accompanying drawing, in which:


Fig. 1 is a flow chart depicting the steps of a method for imaging a thrombus in accordance with the present invention.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention relates to a method for imaging a thrombus, the steps of which are depicted in Fig. 1. At step 10, the patient is administered a radiolabelled compound that preferentially binds to the thrombus. For example, the radiolabelled compound may be administered by injecting approximately 20 mCi (740 Mbq) of the radiolabelled compound into the venous circulation system of the patient. In one embodiment, the radiolabelled compound comprises a radiopharmaceutical of the type described in U.S. Patent No. 5,744,120 issued April 28, 1998 to Edwards et al., U.S. Patent

No. 5,879,657 issued March 9, 1999 to DeGrado et al., U.S. Patent No. 5,879,659 issued March 9, 1999 to Edwards et al., and U.S. Patent No. 5,750,088 issued May 12, 1998 to Sworin et al., all of which are incorporated herein by reference.

Specifically, a radiopharmaceutical useful as an imaging agent in accordance with the present invention is given by formula (I):




wherein d' is preferably between about 1 and 20, x is independently 1-2; y is independently 1-2; and z is independently 0-4.

Q is a glycoprotein IIb/IIIa binding compound selected from the group including the cyclic IIb/IIIa receptor antagonist compounds described in co-pending U.S. Serial No.08/415,908,861 (equivalent to WO 94/22494); the RGD containing peptides described in U.S. Patent Nos. 4,578,079 and 4,792,525, the patent applications PCT US88/04403, PCT US89/01742, PCT US90/03788, and PCT US91/02356, and by Ojima et. al., 204th Meeting of the Amer. Chem. Soc., 1992, Abstract 44; the peptides that are fibrinogen receptor antagonists described in European Patent Application Nos. 90202015.5, 90202030.4, 90202032.2, 90202032.0, 90311148.2, 90311151.6, and 90311537.6; the specific binding peptides and polypeptides described as IIb/IIIa receptor ligands, ligands for the polymerization site of fibrin, laminin derivatives, ligands for fibrinogen, or thrombin ligands in PCT WO 93/23085 (excluding the technetium binding groups); the oligopeptides that correspond to the IIIa protein described in PCT WO 90/00178; the hirudin-based peptides described in PCT WO 90/03391; the IIb/IIIa receptor ligands described in PCT WO 90/15818; the thrombus, platelet or atherosclerotic plaque binding peptides described in PCT WO 92/13572 (excluding the technetium binding group) and GB 9313965.7; the fibrin binding peptides described in U.S. Patent Nos. 4,427,646 and

5,270,030; the hirudin-based peptides described in U.S. Patent No. 5,279,812; the fibrin binding proteins described in U.S. Patent No. 5,217,705; the guanine derivatives that bind to the IIb/IIIa receptor described in U.S. Patent No. 5,086,069; the tyrosine derivatives described in European Patent Application No. 0478328A1, and Hartman et. al., J. Med. Chem., 1992, 35, 4640; or an oxidized low density lipoprotein (LDL).

In one embodiment, Q is of the formula (II):



(II)

or a pharmaceutically acceptable salt or prodrug form thereof wherein:

R³¹ is a C₆-C₁₄ saturated, partially saturated, or aromatic carbocyclic ring system substituted with 0-4 R¹⁰ or R^{10a};

R³² is selected from:

- C(=O)-;
- C(=S)-
- S(=O)₂-;
- S(=O)-;
- P(=Z) (ZR¹³)-;

Z is S or O;

n'' and n' are independently 0-2;

R¹ and R²² are independently selected from the following groups:

hydrogen,

C₁-C₈ alkyl substituted with 0-2 R¹¹;
C₂-C₈ alkenyl substituted with 0-2 R¹¹;
C₂-C₈ alkynyl substituted with 0-2 R¹¹;
C₃-C₁₀ cycloalkyl substituted with 0-2 R¹¹;

aryl substituted with 0-2 R¹²;

a 5-10-membered heterocyclic ring system containing
1-4 heteroatoms independently selected from N, S,
and O, said heterocyclic ring being substituted with
0-2 R¹²;

=O, F, Cl, Br, I, -CF₃, -CN, -CO₂R¹³, -C(=O)R¹³,
-C(=O)N(R¹³)₂, -CHO, -CH₂OR¹³, -OC(=O)R¹³,
-OC(=O)OR^{13a}, -OR¹³, -OC(=O)N(R¹³)₂, -NR¹³C(=O)R¹³,
-NR¹⁴C(=O)OR^{13a}, -NR¹³C(=O)N(R¹³)₂, -NR¹⁴SO₂N(R¹³)₂,
-NR¹⁴SO₂R^{13a}, -SO₃H, -SO₂R^{13a}, -SR¹³, -S(=O)R^{13a},
-SO₂N(R¹³)₂, -N(R¹³)₂, -NHC(=NH)NHR¹³, -C(=NH)NHR¹³,
=NOR¹³, NO₂, -C(=O)NHR¹³, -C(=O)NHN(R¹³)R^{13a},
-OCH₂CO₂H, 2-(1-morpholino)ethoxy;

R¹ and R²¹ can alternatively join to form a 3-7 membered
carbocyclic ring substituted with 0-2 R¹²;

when n' is 2, R¹ or R²¹ can alternatively be taken
together with R¹ or R²¹ on an adjacent carbon atom to
form a direct bond, thereby to form a double or triple
bond between said carbon atoms;

R²² and R²³ can alternatively join to form a 3-7 membered
carbocyclic ring substituted with 0-2 R¹²;

when n" is 2, R²² or R²³ can alternatively be taken

together with R²² or R²³ on an adjacent carbon atom to form a direct bond, thereby to form a double or triple bond between the adjacent carbon atoms;

R¹ and R², where R²¹ is H, can alternatively join to form a 5-8 membered carbocyclic ring substituted with 0-2 R¹²;

R¹¹ is selected from one or more of the following:

=O, F, Cl, Br, I, -CF₃, -CN, -CO₂R¹³, -C(=O)R¹³,
-C(=O)N(R¹³)₂, -CHO, -CH₂OR¹³, -OC(=O)R¹³,
-OC(=O)OR^{13a}, -OR¹³, -OC(=O)N(R¹³)₂, -NR¹³C(=O)R¹³,
-NR¹⁴C(=O)OR^{13a}, -NR¹³C(=O)N(R¹³)₂, -NR¹⁴SO₂N(R¹³)₂,
-NR¹⁴SO₂R^{13a}, -SO₃H, -SO₂R^{13a}, -SR¹³, -S(=O)R^{13a},
-SO₂N(R¹³)₂, -N(R¹³)₂, -NHC(=NH)NHR¹³, -C(=NH)NHR¹³,
=NOR¹³, NO₂, -C(=O)NHOR¹³, -C(=O)NHN(R¹³)R^{13a},
-OCH₂CO₂H, 2-(1-morpholino)ethoxy,

C₁-C₅ alkyl, C₂-C₄ alkenyl, C₃-C₆ cycloalkyl, C₃-C₆
cycloalkylmethyl, C₂-C₆ alkoxyalkyl, C₃-C₆
cycloalkoxy, C₁-C₄ alkyl (alkyl being substituted
with 1-5 groups selected independently from:
-NR¹³R¹⁴, -CF₃, NO₂, -SO₂R^{13a}, or -S(=O)R^{13a}),

aryl substituted with 0-2 R¹²,

a 5-10-membered heterocyclic ring system containing
1-4 heteroatoms independently selected from N, S,
and O, said heterocyclic ring being substituted with
0-2 R¹²;

R¹² is selected from one or more of the following:

phenyl, benzyl, phenethyl, phenoxy, benzyloxy,
halogen, hydroxy, nitro, cyano, C₁-C₅ alkyl, C₃-C₆
cycloalkyl, C₃-C₆ cycloalkylmethyl, C₇-C₁₀
arylalkyl, C₁-C₅ alkoxy, -CO₂R¹³, -C(=O)NHOR^{13a},

-C(=O)NHN(R¹³)₂, =NOR¹³, -B(R³⁴)(R³⁵), C₃-C₆
cycloalkoxy, -OC(=O)R¹³, -C(=O)R¹³, -OC(=O)OR^{13a},
-OR¹³, -(C₁-C₄ alkyl)-OR¹³, -N(R¹³)₂,
-OC(=O)N(R¹³)₂, -NR¹³C(=O)R¹³, -NR¹³C(=O)OR^{13a},
-NR¹³C(=O)N(R¹³)₂, -NR¹³SO₂N(R¹³)₂, -NR¹³SO₂R^{13a},
-SO₃H, -SO₂R^{13a}, -S(=O)R^{13a}, -SR¹³, -SO₂N(R¹³)₂,
C₂-C₆ alkoxyalkyl, methylenedioxy, ethylenedioxy,
C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, C₁-C₄
alkylcarbonyloxy, C₁-C₄ alkylcarbonyl, C₁-C₄
alkylcarbonylamino, -OCH₂CO₂H,
2-(1-morpholino)ethoxy, C₁-C₄ alkyl (alkyl being
substituted with -N(R¹³)₂, -CF₃, NO₂, or
-S(=O)R^{13a});

R¹³ is selected independently from: H, C₁-C₁₀ alkyl,
C₃-C₁₀ cycloalkyl, C₄-C₁₂ alkylcycloalkyl, aryl, -(C₁-C₁₀
alkyl)aryl, or C₃-C₁₀ alkoxyalkyl;

R^{13a} is C₁-C₁₀ alkyl, C₃-C₁₀ cycloalkyl, C₄-C₁₂
alkylcycloalkyl, aryl, -(C₁-C₁₀ alkyl)aryl, or C₃-C₁₀
alkoxyalkyl;

when two R¹³ groups are bonded to a single N, said R¹³
groups may alternatively be taken together to form
-(CH₂)₂₋₅- or -(CH₂)O(CH₂)-;

R¹⁴ is OH, H, C₁-C₄ alkyl, or benzyl;

R²¹ and R²³ are independently selected from:

hydrogen;

C₁-C₄ alkyl, optionally substituted with 1-6
halogen;

benzyl;

R² is H or C₁-C₈ alkyl;

R¹⁰ and R^{10a} are selected independently from one or more of the following:

phenyl, benzyl, phenethyl, phenoxy, benzyloxy,
halogen, hydroxy, nitro, cyano, C₁-C₅ alkyl, C₃-C₆
cycloalkyl, C₃-C₆ cycloalkylmethyl, C₇-C₁₀
arylalkyl, C₁-C₅ alkoxy, -CO₂R¹³, -C(=O)N(R¹³)₂,
-C(=O)NHOR^{13a}, -C(=O)NHN(R¹³)₂, =NOR¹³,
-B(R³⁴)(R³⁵), C₃-C₆ cycloalkoxy, -OC(=O)R¹³,
-C(=O)R¹³, -OC(=O)OR^{13a}, -OR¹³, -(C₁-C₄ alkyl)-OR¹³,
-N(R¹³)₂, -OC(=O)N(R¹³)₂, -NR¹³C(=O)R¹³,
-NR¹³C(=O)OR^{13a}, -NR¹³C(=O)N(R¹³)₂, -NR¹³SO₂N(R¹³)₂,
-NR¹³SO₂R^{13a}, -SO₃H, -SO₂R^{13a}, -S(=O)R^{13a}, -SR¹³,
-SO₂N(R¹³)₂, C₂-C₆ alkoxyalkyl, methylenedioxy,
ethylenedioxy, C₁-C₄ haloalkyl (including -C_vF_w
where v = 1 to 3 and w = 1 to (2v+1)), C₁-C₄
haloalkoxy, C₁-C₄ alkylcarbonyloxy, C₁-C₄
alkylcarbonyl, C₁-C₄ alkylcarbonylamino, -OCH₂CO₂H,
2-(1-morpholino)ethoxy, C₁-C₄ alkyl (alkyl being
substituted with -N(R¹³)₂, -CF₃, NO₂, or
-S(=O)R^{13a});

J is 3-aminopropionic acid or an L-isomer or D-isomer
amino acid of structure -N(R³)C(R⁴)(R⁵)C(=O)-, wherein:

R³ is H or C₁-C₈ alkyl;

R⁴ is H or C₁-C₃ alkyl;

R⁵ is selected from:

hydrogen;

C₁-C₈ alkyl substituted with 0-2 R¹¹;

C₂-C₈ alkenyl substituted with 0-2 R¹¹;

C₂-C₈ alkynyl substituted with 0-2 R¹¹;


C₃-C₁₀ cycloalkyl substituted with 0-2 R¹¹;

aryl substituted with 0-2 R¹²;

a 5-10-membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, or O, said heterocyclic ring being substituted with 0-2 R¹²;


=O, F, Cl, Br, I, -CF₃, -CN, -CO₂R¹³, -C(=O)R¹³, -C(=O)N(R¹³)₂, -CHO, -CH₂OR¹³, -OC(=O)R¹³, -OC(=O)OR^{13a}, -OR¹³, -OC(=O)N(R¹³)₂, -NR¹³C(=O)R¹³, -NR¹⁴C(=O)OR^{13a}, -NR¹³C(=O)N(R¹³)₂, -NR¹⁴SO₂N(R¹³)₂, -NR¹⁴SO₂R^{13a}, -SO₃H, -SO₂R^{13a}, -SR¹³, -S(=O)R^{13a}, -SO₂N(R¹³)₂, -N(R¹³)₂, -NHC(=NH)NHR¹³, -C(=NH)NHR¹³, =NOR¹³, NO₂, -C(=O)NHOR¹³, -C(=O)NHN(R¹³)R^{13a}, =NOR¹³, -B(R³⁴)(R³⁵), -OCH₂CO₂H, 2-(1-morpholino)ethoxy, -SC(=NH)NHR¹³, N₃, -Si(CH₃)₃, (C₁-C₅ alkyl)NHR¹⁶;

-(C₀-C₆ alkyl)X;



, where q is

independently 0,1;




;

-(CH₂)_mS(O)_p(CH₂)₂X, where m = 1,2 and p' = 0-2;

wherein X is defined below; and

R³ and R⁴ may also be taken together to form



;

R³ and R⁵ can alternatively be taken together to form
-(CH₂)_t- or -CH₂S(O)_p'C(CH₃)₂-, where t = 2-4 and p' =
0-2; or

R⁴ and R⁵ can alternatively be taken together to form
-(CH₂)_u-, where u = 2-5;

R¹⁶ is selected from:

an amine protecting group;

1-2 amino acids;

1-2 amino acids substituted with an amine protecting
group;


K is a D-isomer or L-isomer amino acid of structure

-(R⁶)CH(R⁷)C(=O)-, wherein:

R⁶ is H or C₁-C₈ alkyl;


R⁷ is selected from:

-(C₁-C₇ alkyl)X;




, wherein each q is

independently 0-2 and substitution on the phenyl is at the 3 or 4 position;



, wherein each


q is independently 0-2 and substitution on the cyclohexyl is at the 3 or 4 position;



$\text{---}(\text{CH}_2)_m\text{O---}(\text{C}_1\text{---C}_4\text{ alkyl})\text{---X}$, where m = 1 or 2;

$\text{---}(\text{CH}_2)_m\text{S(O)}\text{P'---}(\text{C}_1\text{---C}_4\text{ alkyl})\text{---X}$, where m = 1 or 2 and p' = 0-2; and


X is selected from:




;

-N(R¹³)R¹³; -C(=NH)(NH₂); -SC(=NH)-NH₂; -NH-C(=NH)(NHCN);
 -NH-C(=NCN)(NH₂); -NH-C(=N-OR¹³)(NH₂);

R⁶ and R⁷ can alternatively be taken together to form




or 2 and wherein n = 0 or 1 and X is -NH₂ or



L is -Y(CH₂)_vC(=O)-, wherein:

Y is NH, N(C₁-C₃ alkyl), O, or S; and v = 1 or 2;

M is a D-isomer or L-isomer amino acid of structure



, wherein:

q' is 0-2;

R¹⁷ is H, C₁-C₃ alkyl;

R⁸ is selected from:

-CO₂R¹³, -SO₃R¹³, -SO₂NHR¹⁴, -B(R³⁴)(R³⁵), -NHSO₂CF₃,
-CONHNHSO₂CF₃, -PO(OR¹³)₂, -PO(OR¹³)R¹³,
-SO₂NH-heteroaryl (said heteroaryl being
5-10-membered and having 1-4 heteroatoms selected
independently from N, S, or O), -SO₂NH-heteroaryl
(said heteroaryl being 5-10-membered and having 1-4
heteroatoms selected independently from N, S, or O),
-SO₂NHCOR¹³, -CONHSO₂R^{13a}, -CH₂CONHSO₂R^{13a},
-NHSO₂NHCOR^{13a}, -NHCONHSO₂R^{13a}, -SO₂NHCONHR¹³;

R³⁴ and R³⁵ are independently selected from:

-OH,
-F,
-N(R¹³)₂, or
C₁-C₈-alkoxy;


R³⁴ and R³⁵ can alternatively be taken together form:

a cyclic boron ester where said chain or ring
contains from 2 to 20 carbon atoms and, optionally,
1-4 heteroatoms independently selected from N, S, or
O;

a divalent cyclic boron amide where said chain or
ring contains from 2 to 20 carbon atoms and,
optionally, 1-4 heteroatoms independently selected
from N, S, or O;

a cyclic boron amide-ester where said chain or ring
contains from 2 to 20 carbon atoms and, optionally,
1-4 heteroatoms independently selected from N, S, or
O.

In another embodiment, Q is of the formula (III):



(III)

or a pharmaceutically acceptable salt or prodrug form thereof wherein:

the shown phenyl ring may be further substituted with 0-3 R¹⁰;

R¹⁰ is selected independently from: H, C₁-C₈ alkyl, phenyl, halogen, or C₁-C₄ alkoxy;

R¹ is H, C₁-C₄ alkyl, phenyl, benzyl, or phenyl-(C₁-C₄)alkyl;

R² is H or methyl;

R¹³ is selected independently from: H, C₁-C₁₀ alkyl, C₃-C₁₀ cycloalkyl, C₄-C₁₂ alkylcycloalkyl, aryl, -(C₁-C₁₀ alkyl)aryl, or C₃-C₁₀ alkoxyalkyl;

R¹³ᵃ is C₁-C₁₀ alkyl, C₃-C₁₀ cycloalkyl, C₄-C₁₂ alkylcycloalkyl, aryl, -(C₁-C₁₀ alkyl)aryl, or C₃-C₁₀ alkoxyalkyl;

when two R^{13} groups are bonded to a single N, said R^{13} groups may alternatively be taken together to form $-(CH_2)_2-$ or $-(CH_2)O(CH_2)-$;

R^{14} is OH, H, C_1 - C_4 alkyl, or benzyl;

J is β -alanine or an L-isomer or D-isomer amino acid of structure $-N(R^3)C(R^4)(R^5)C(=O)-$, wherein:

R^3 is H or CH_3 ;

R^4 is H or C_1 - C_3 alkyl;

R^5 is H, C_1 - C_8 alkyl, C_3 - C_6 cycloalkyl, C_3 - C_6 cycloalkylmethyl, C_1 - C_6 cycloalkylethyl, phenyl, phenylmethyl, CH_2OH , CH_2SH , CH_2OCH_3 , CH_2SCH_3 , $CH_2CH_2SCH_3$, $(CH_2)_sNH_2$, $-(CH_2)_sNHC(=NH)(NH_2)$, $-(CH_2)_sNHR^{16}$, where $s = 3-5$; or

R^{16} is selected from:

an amine protecting group;

1-2 amino acids; or

1-2 amino acids substituted with an amine protecting group;


R^3 and R^5 can alternatively be taken together to form $-CH_2CH_2CH_2-$; or

R^4 and R^5 can alternatively be taken together to form $-(CH_2)_u-$, where $u = 2-5$;

K is an L-isomer amino acid of structure $-N(R^6)CH(R^7)C(=O)-$, wherein:


R^6 is H or C_1 - C_8 alkyl;

R^7 is:




;


5



$-(CH_2)_rX$, where $r = 3-6$;



;




;

$-(CH_2)_mS(CH_2)_2X$, where $m = 1 \text{ or } 2$;

15

$-(C_3-C_7 \text{ alkyl})-\text{NH}-(C_1-C_6 \text{ alkyl})$;




$\text{---}(\text{CH}_2)_m\text{---O---}(\text{C}_1\text{---C}_4 \text{ alkyl})\text{---NH---}(\text{C}_1\text{---C}_6 \text{ alkyl})$, where $m = 1$ or 2 ;

$\text{---}(\text{CH}_2)_m\text{---S---}(\text{C}_1\text{---C}_4 \text{ alkyl})\text{---NH---}(\text{C}_1\text{---C}_6 \text{ alkyl})$, where $m = 1$ or 2 ; and

X is ---NH_2 or $\text{---NHC(=NH)(NH}_2\text{)}$, provided that X is not ---NH_2 when $r = 4$; or

R^6 and R^7 are alternatively be taken together to form




$\text{---NHC(=NH)(NH}_2\text{)}$;

L is $\text{---Y(CH}_2\text{)}_v\text{C(=O)---}$, wherein:

Y is NH, O, or S; and $v = 1, 2$;

M is a D-isomer or L-isomer amino acid of structure



, wherein:

q' is 0-2;

R¹⁷ is H, C₁-C₃ alkyl;

R⁸ is selected from:

-CO₂R¹³, -SO₃R¹³, -SO₂NHR¹⁴, -B(R³⁴)(R³⁵), -NHSO₂CF₃,
-CONHNHSO₂CF₃, -PO(OR¹³)₂, -PO(OR¹³)R¹³,
-SO₂NH-heteroaryl (said heteroaryl being
5-10-membered and having 1-4 heteroatoms selected
independently from N, S, or O), -SO₂NH-heteroaryl
(said heteroaryl being 5-10-membered and having 1-4
heteroatoms selected independently from N, S, or O),
-SO₂NHCOR¹³, -CONHSO₂R^{13a}, -CH₂CONHSO₂R^{13a},
-NHSO₂NHCOR^{13a}, -NHCONHSO₂R^{13a}, -SO₂NHCONHR¹³.

C_n is a radionuclide metal chelator or bonding unit bound to the biologically active compound Q, either directly or through the optional linking group L_n. C_n is preferably selected from the group consisting of: R⁴⁰N=N⁺=, R⁴⁰R⁴¹N=N-, R⁴⁰N=, and R⁴⁰N=N(H)-; wherein,

R⁴⁰ is independently selected at each occurrence from the group consisting of: a bond to L_n, C₁-C₁₀ alkyl substituted with 0-3 R⁵², aryl substituted with 0-3 R⁵², cycloalkyl substituted with 0-3 R⁵², heterocycle substituted with 0-3 R⁵², heterocycloalkyl substituted with 0-3 R⁵², aralkyl substituted with 0-3 R⁵² and alkaryl substituted with 0-3 R⁵²;

R⁴¹ is independently selected from the group consisting of: hydrogen, aryl substituted with 0-3 R⁵², C₁-C₁₀ alkyl substituted with 0-3 R⁵², and a heterocycle substituted with 0-3 R⁵²;


R⁵² is independently selected at each occurrence from the group consisting of: a bond to L_n, =O, F, Cl, Br, I, -CF₃, -CN, -CO₂R⁵³, -C(=O)R⁵³, -C(=O)N(R⁵³)₂, -CHO, -CH₂OR⁵³, -OC(=O)R⁵³, -OC(=O)OR^{53a}, -OR⁵³,

$-\text{OC}(=\text{O})\text{N}(\text{R}^{53})_2$, $-\text{NR}^{53}\text{C}(=\text{O})\text{R}^{53}$, $-\text{NR}^{54}\text{C}(=\text{O})\text{OR}^{53a}$,
 $-\text{NR}^{53}\text{C}(=\text{O})\text{N}(\text{R}^{53})_2$, $-\text{NR}^{54}\text{SO}_2\text{N}(\text{R}^{53})_2$, $-\text{NR}^{54}\text{SO}_2\text{R}^{53a}$, $-\text{SO}_3\text{H}$,
 $-\text{SO}_2\text{R}^{53a}$, $-\text{SR}^{53}$, $-\text{S}(=\text{O})\text{R}^{53a}$, $-\text{SO}_2\text{N}(\text{R}^{53})_2$, $-\text{N}(\text{R}^{53})_2$,
 $-\text{NHC}(=\text{NH})\text{NHR}^{53}$, $-\text{C}(=\text{NH})\text{NHR}^{53}$, $-\text{NOR}^{53}$, NO_2 , $-\text{C}(=\text{O})\text{NHR}^{53}$,
 $-\text{C}(=\text{O})\text{NHNHR}^{53}\text{R}^{53a}$, $-\text{OCH}_2\text{CO}_2\text{H}$, 2-(1-morpholino)ethoxy; and

R^{53} , R^{53a} , and R^{54} are each independently selected at
 each occurrence from the group consisting of: hydrogen,
 $\text{C}_1\text{-C}_6$ alkyl, and a bond to L_n .

In order to have a chelating diazenido group (i.e., a group of
 formula $\text{R}^{40}\text{N}=\text{N}'$ or $\text{R}^{40}\text{N}=(\text{H})-$ at least one other atom of the
 group located on R^{40} must also be bound to the radionuclide.
 The atoms bound to the metal are termed donor atoms.

The optional linking group L_n is given by the formula:



wherein:

M^1 is $-\{(\text{CH}_2)_g \text{Z}^1\}_{g'} - (\text{CR}^{55}\text{R}^{56})_{g''} -$;

M^2 is $-(\text{CR}^{55}\text{R}^{56})_{g''} - [\text{Z}^1 (\text{CH}_2)_g]_{g'} -$;

g is independently 0-10;

g' is independently 0-1;

g'' is independently 0-10;

f is independently 0-10;

f' is independently 0-10;

f'' is independently 0-1;

Y^1 and Y^2 , are independently selected at each

occurrence from: a bond, O, NR⁵⁶, C=O, C(=O)O, OC(=O)O, C(=O)NH-, C=NR⁵⁶, S, SO, SO₂, SO₃, NHC(=O), (NH)₂C(=O), and (NH)₂C=S;

Z¹ is independently selected at each occurrence from a C₆-C₁₄ saturated, partially saturated, or aromatic carbocyclic ring system, substituted with 0-4 R⁵⁷; and a heterocyclic ring system, substituted with 0-4 R⁵⁷;

R⁵⁵ and R⁵⁶ are independently selected at each occurrence from the group consisting of: hydrogen; C₁-C₁₀ alkyl substituted with 0-5 R⁵⁷; and alkaryl wherein the aryl is substituted with 0-5 R⁵⁷;

R⁵⁷ is independently selected at each occurrence from the group: hydrogen, OH, NHR⁵⁸, C(=O)R⁵⁸, OC(=O)R⁵⁸, OC(=O)OR⁵⁸, C(=O)OR⁵⁸, C(=O)NR⁵⁸, C=N, SR⁵⁸, SOR⁵⁸, SO₂R⁵⁸, NHC(=O)R⁵⁸, NHC(=O)NHR⁵⁸, NHC(=S)NHR⁵⁸; or, alternatively, when attached to an additional molecule Q, R⁵⁷ is independently selected at each occurrence from the group: O, NR⁵⁸, C=O, C(=O)O, OC(=O)O, C(=O)N-, C=NR⁵⁸, S, SO, SO₂, SO₃, NHC(=O), (NH)₂C(=O), (NH)₂C=S; and

R⁵⁸ is independently selected at each occurrence from the group: hydrogen; C₁-C₆ alkyl; benzyl, and phenyl.

The radiopharmaceutical compound used in accordance with the present invention is radiolabelled. By "radiolabelled", it is meant that the compound contains a radioisotope which is suitable for administration to a mammalian patient. Suitable radioisotopes are known to those skilled in the art and include, for example, isotopes of halogens (such as chlorine,

fluorine, bromine and iodine), technetium and indium. Preferable radioisotopes include ^{123}I , ^{125}I , ^{131}I , $^{99\text{m}}\text{Tc}$, and ^{111}In , more preferably ^{111}In , ^{123}I and $^{99\text{m}}\text{Tc}$, and most preferably $^{99\text{m}}\text{Tc}$. Radiolabelled compounds of the invention may be prepared using standard radiolabelling procedures well known to those skilled in the art. The glycoprotein IIB/IIIA binding compound, Q, is radiolabelled indirectly (that is, by incorporating the radiolabel into the compound through the chelating agent C_h). Such radiolabelling should also be reasonably stable, both chemically and metabolically, applying recognized standards in the art. Also, although the radiolabelled compound may be labeled in a variety of fashions with a variety of different radioisotopes, as those skilled in the art will recognize, such radiolabelling should be carried out in a manner such that the high binding affinity and specificity of the unlabeled glycoprotein IIB/IIIA binding compound to the glycoprotein IIB/IIIA receptor is not significantly affected. By not significantly affected, it is meant that the binding affinity and specificity is not affected more than about 50%, preferably not more than about 40%, more preferably not more than about 30%, even more preferably not more than about 20%, and still even more preferably not more than about 10%, and most preferably the binding affinity and specificity is not affected at all.

Referring again to formula (I), M_r is a transition metal radionuclide which is attached to the biologically active compound Q via the chelator C_h . Preferred radiolabelled compounds of the invention are radiolabelled compounds wherein the radiolabel is located on the carbocyclic ring system of R^{31} of formula (II). Even more preferred radiolabelled compounds of the invention are those of formula (III), wherein the radiolabel is located at position R^{10} or R^{10a} substituted on the benzene ring.

The coordination sphere of the radionuclide includes all the ligands or groups bound to the radionuclide. For a transition metal radionuclide, M_t , to be stable it typically has a coordination number comprised of an integer greater than

or equal to 5 and less than or equal to 7; that is there are 5 to 7 atoms bound to the metal and it is said to have a complete coordination sphere. If the chelator or bonding unit Ch' does not provide all of the atoms necessary to stabilize the metal radionuclide by completing its coordination sphere, the coordination sphere is completed by donor atoms from other ligands, termed ancillary or co-ligands, which can be either terminal or chelating.

A large number of ligands can serve as ancillary or co-ligands, the choice of which is determined by a variety of considerations such as the ease of synthesis of the radiopharmaceutical, the chemical and physical properties of the ancillary ligand, the rate of formation, the yield, the number of isomeric forms of the resulting radiopharmaceuticals, the ability to administer said ancillary or co-ligand to a patient without adverse physiological consequences to said patient, and the compatibility of the ligand in a lyophilized kit formulation. The charge and lipophilicity of the ancillary ligand will effect the charge and lipophilicity of the radiopharmaceuticals. For example, the use of 4,5-dihydroxy-1,3-benzene disulfonate results in radiopharmaceuticals with an additional two anionic groups because the sulfonate groups will be anionic under physiological conditions. The use of N-alkyl substituted 3,4-hydroxypyridinones results in radiopharmaceuticals with varying degrees of lipophilicity depending on the size of the alkyl substituents.

The radiopharmaceuticals prepared from the reagents of the present invention can be comprised of one or two ancillary or co-ligands, designated Al_1 , in a binary ligand system. The one or two ancillary or co-ligands, Al_1 , comprising the radiopharmaceuticals can be independently selected from the group consisting of: dioxygen ligands, functionalized aminocarboxylates and halides; provided that the coordination sphere of the radionuclide is complete.

Ancillary dioxygen ligands include ligands that coordinate to the metal ion through at least two oxygen donor

atoms. Examples include but are not limited to:
glucoheptonate, gluconate, 2-hydroxyisobutyrate, lactate,
tartrate, mannitol, glucarate, maltol, Kojic acid,
2,2-bis(hydroxymethyl)propionic acid,
4,5-dihydroxy-1,3-benzene disulfonate, or substituted or
unsubstituted 1,2 or 3,4 hydroxypyridinones, or
pharmaceutically acceptable salts thereof.


Functionalized aminocarboxylates include ligands that
have a combination of nitrogen and oxygen donor atoms.
Examples include but are not limited to: iminodiacetic acid,
2,3 diaminopropionic acid, nitrilotriacetic acid,
N,N'-ethylenediamine diacetic acid, N,N,N'-ethylenediamine
triacetic acid, hydroxyethylethylenediamine triacetic acid,
N,N'-ethylenediamine bis-hydroxyphenylglycine, or the ligands
described in Eur. Pat. Appl. No. 93302712.0, or
pharmaceutically acceptable salts thereof.

Halides which are suitable for use as the ancillary
ligand AL_1 can be chloride, bromide, fluoride, iodide, or
pharmaceutically acceptable salts thereof.

Of particular utility are radiopharmaceuticals prepared
from the reagents of the present invention comprised of two
different types of ancillary or co-ligands, one or two ligands
designated the first ancillary or co-ligand or ligands, AL_1 ,
and independently selected from the group: dioxygen ligands,
functionalized aminocarboxylates and halides; and one to four
ligands designated the second ancillary or co-ligand or
ligands, AL_2 , selected from the group: trisubstituted
phosphines, trisubstituted arsines, tetrasubstituted
diphosphines and tetrasubstituted diarsines, in a ternary
ligand system. Radiopharmaceuticals comprised of one or more
ancillary or co-ligands AL_2 are more stable compared to said
radiopharmaceuticals that are not comprised of one or more
ancillary ligands, AL_2 ; that is, they have a minimal number of
isomeric forms, the relative ratios of which do not change
significantly with time, and remain substantially intact upon
dilution.


In one particular embodiment, the radiopharmaceutical

comprises a compound of the formula (IV):



(IV) .

In another particular embodiment, the radiopharmaceutical comprises a compound of the formula (V):



(V) .

After the radiopharmaceutical has been administered, a series of image slices of the thrombus are acquired at step 12 of Fig. 1. The image slices reflect the concentration of radioactivity within the thrombus. Each image slice is composed of a two-dimensional array of pixels, wherein each pixel comprises an intensity value representative of the concentration of radioactivity at the particular position within the thrombus which corresponds to the pixel. In one embodiment, the image slices are obtained using a gamma camera to record single photon emission computed tomography (SPECT) images.

When the method of the present invention is used to detect an arterial thrombus, for example, data should be acquired using parameters that enhance the sensitivity of the technique for small lesions. In particular, when SPECT images are acquired, the parameters should include a 3 mm or smaller digital sampling (i.e., pixel size), and a minimum of 90 views over a 360 degree rotation or 45 views over a 180 degree rotation. Further, very high-resolution collimators and multiple heads should be used to increase the resolution and sensitivity, respectively. In addition, the acquired data should be reconstructed using a spatial filter with a relatively high frequency cutoff (i.e., approximately 3.00 or higher) to minimize resolution loss due to smoothing. A minimum of thirty angles ensures a visually smooth effect when the views are intended to be displayed sequentially to produce a rotating view of the thrombus, as described below.

At step 14, the image slices are reconstructed and assembled into a three-dimensional matrix of data. The three-dimensional matrix of data is assembled by stacking the individual image slices in sequential order. When the image slices are collected perpendicularly to the long axis of the patient, the three-dimensional matrix of data is organized as a series of transaxial image slices.

Once the three-dimensional matrix of data has been assembled, the matrix of data is scanned along an array of parallel lines (usually perpendicular to the vertical axis of

the body), at step 16, to determine the maximum intensity value along each of the parallel lines. The maximum intensity value along a parallel line is equivalent to the most intense pixel within the matrix of data encountered along that parallel line. The most intense pixel is defined as the pixel corresponding to the position within the lesion where the radioactivity is most intense.

At step 18, the maximum value along each parallel line is assigned to a pixel in a two-dimensional image array. The relative position of the pixel in the two-dimensional image array corresponds to the relative position of the line in the array of parallel lines. The resulting two-dimensional image array therefore represents an image of the most intense pixels as viewed by an observer viewing the thrombus along the array of parallel lines.

The three-dimensional matrix of data can be scanned along additional arrays of parallel lines in order to produce image arrays of the thrombus from different angles. At step 20, it is determined whether the three-dimensional matrix of data is to be scanned at a different angle. If the matrix of data is to be scanned at a different angle, the matrix of data is scanned along the new angle at step 16. Preferably, the lesion is scanned along a minimum of 90 views over a series of angles over 360° or a minimum of 45 views over a series of angles over 180°.

If additional views of the thrombus are not desired, the results are displayed at step 22. The results can be displayed as individual views of the thrombus by displaying one or more of the individual two-dimensional image arrays. Alternatively, when two-dimensional image arrays at more than one angle have been obtained, the individual two-dimensional image arrays can be displayed sequentially by angle to produce a rotating view of the most intense pixels.


Although the above discussion has focused primarily on localization of pulmonary embolism, the present invention is not intended to be so limited. Instead, the present invention is intended to relate to any medical condition capable of

diagnosis using a clot-binding radiopharmaceutical contrast agent and SPECT. For example, it is recognized that the present invention is equally applicable to localization of thrombii in general and arterial coronary thrombii in particular.

It will be recognized by those skilled in the art that changes or modifications may be made to the above-described embodiments without departing from the broad inventive concepts of the invention. It should therefore be understood that this invention is not limited to the particular embodiments described herein, but is intended to include all changes and modifications that are within the scope and spirit of the invention as set forth in the claims.

WHAT IS CLAIMED IS:

1. A method for imaging a thrombus comprising the steps of:
 - a. localizing a radiolabelled compound at the thrombus;
 - b. acquiring image slices representing a physical property of the radiolabelled thrombus;
 - c. assembling the image slices into a three-dimensional matrix of data;
 - d. scanning the three-dimensional matrix of data along an array of parallel lines to determine a maximum value along each line; and
 - e. assigning the maximum value along each line to a pixel in a two-dimensional array, the position of the pixel corresponding to the position of the line in the array of parallel lines.
2. The method of Claim 1 wherein the localization step comprises the step of localizing a compound that preferentially binds to activated platelets of the thrombus.
3. The method of Claim 2 wherein the localization step comprises the step of localizing a compound that binds to activated platelets of the thrombus via the glycoprotein IIb/IIIa receptor.
4. The method of Claim 3 wherein the localization step comprises the step of localizing a compound of the formula (I), and pharmaceutically acceptable salts thereof, at the thrombus:




(I),

wherein,

Q is a glycoprotein IIb/IIIa binding compound;

d' is 1 - 20;

L_n is a linking group of formula:



wherein:

M¹ is - [(CH₂)_g Z¹]_{g'} - (CR⁵⁵R⁵⁶)_{g''} -;

M² is - (CR⁵⁵R⁵⁶)_{g''} - [Z¹ (CH₂)_g]_{g'} -;

g is independently 0-10;

g' is independently 0-1;

g'' is independently 0-10;

f is independently 0-10;

f' is independently 0-10;

f'' is independently 0-1;

Y¹ and Y², are independently selected at each occurrence from: a bond, O, NR⁵⁶, C=O, C(=O)O, OC(=O)O, C(=O)NH-, C=NR⁵⁶, S, SO, SO₂, SO₃, NHC(=O), (NH)₂C(=O), and (NH)₂C=S;

Z¹ is independently selected at each occurrence from a C₆-C₁₄ saturated, partially saturated, or aromatic carbocyclic ring system, substituted with 0-4 R⁵⁷; and a heterocyclic ring system, substituted with 0-4 R⁵⁷;

R⁵⁵ and R⁵⁶ are independently selected at each occurrence from: hydrogen; C₁-C₁₀ alkyl substituted with 0-5 R⁵⁷; and alkaryl wherein the aryl is substituted with 0-5 R⁵⁷;

R⁵⁷ is independently selected at each occurrence from the group: hydrogen, OH, NHR⁵⁸, C(=O)R⁵⁸, OC(=O)R⁵⁸, OC(=O)OR⁵⁸, C(=O)OR⁵⁸, C(=O)NR⁵⁸, C=N, SR⁵⁸, SOR⁵⁸, SO₂R⁵⁸, NHC(=O)R⁵⁸, NHC(=O)NHR⁵⁸, NHC(=S)NHR⁵⁸; or, alternatively, when attached to an additional molecule Q, R⁵⁷ is independently selected at each occurrence from the group: O, NR⁵⁸, C=O, C(=O)O, OC(=O)O, C(=O)N-, C=NR⁵⁸, S, SO, SO₂, SO₃, NHC(=O), (NH)₂C(=O), (NH)₂C=S; and,

R⁵⁸ is independently selected at each occurrence from the group: hydrogen; C₁-C₆ alkyl; benzyl, and phenyl;

M_r is a transition metal radionuclide;

C_n is a radionuclide metal chelator or bonding unit bound to the transition metal radionuclide selected from the group consisting of: R⁴⁰N=N⁺=, R⁴⁰R⁴¹N-N=, R⁴⁰N=, or R⁴⁰N=N(H)-;

R⁴⁰ is independently selected at each occurrence from the group: a bond to L_n, C₁-C₁₀ alkyl substituted with 0-3 R⁵², aryl substituted with 0-3 R⁵², cycloalkyl substituted with 0-3 R⁵², heterocycle substituted with 0-3 R⁵², heterocycloalkyl substituted with 0-3 R⁵², aralkyl substituted with 0-3 R⁵² and alkaryl substituted with 0-3 R⁵²;

R⁴¹ is independently selected from the group: hydrogen, aryl substituted with 0-3 R⁵², C₁-C₁₀ alkyl substituted with 0-3 R⁵², and a heterocycle substituted with 0-3 R⁵²;

R⁵² is independently selected at each occurrence from the group: a bond to L_n, =O, F, Cl, Br, I, -CF₃, -CN, -CO₂R⁵³, -C(=O)R⁵³, -C(=O)N(R⁵³)₂, -CHO, -CH₂OR⁵³, -OC(=O)R⁵³, -OC(=O)OR^{53a}, -OR⁵³, -OC(=O)N(R⁵³)₂, -NR⁵³C(=O)R⁵³, -NR⁵⁴C(=O)OR^{53a}, -NR⁵³C(=O)N(R⁵³)₂, -NR⁵⁴SO₂N(R⁵³)₂, -NR⁵⁴SO₂R^{53a}, -SO₃H, -SO₂R^{53a}, -SR⁵³, -S(=O)R^{53a}, -SO₂N(R⁵³)₂, -N(R⁵³)₂, -NHC(=NH)NHR⁵³, -C(=NH)NHR⁵³, =NOR⁵³, NO₂, -C(=O)NHR⁵³, -C(=O)NHN(R⁵³)R^{53a}, -OCH₂CO₂H, 2-(1-morpholino)ethoxy;

R⁵³, R^{53a}, and R⁵⁴ are each independently selected at each occurrence from the group: hydrogen, C₁-C₆ alkyl, and a bond to L_n;

A_{L1} is a first ligand wherein each of the y first ligands are selected from the group consisting of: dioxygen ligands, functionalized aminocarboxylates, halides, and combinations thereof;

A_{L2} is a second ligand wherein each of the z second ligands are selected from the group consisting of: trisubstituted phosphines, trisubstituted arsines, tetrasubstituted diphosphines, tetrasubstituted diarsines, and combinations thereof;


x is independently 1-2;

y is independently 1-2; and

z is independently 0-4.

5. The method of Claim 4 wherein M_r is selected from the group consisting of: technetium-99m, rhenium-186, and rhenium-188.
6. The method of Claim 4 wherein the localization step comprises the step of localizing a compound of the

formula (I) at the thrombus wherein Q is of the formula (II),



(II)

or a pharmaceutically acceptable salt or prodrug form thereof wherein:

R³¹ is a C₆-C₁₄ saturated, partially saturated, or aromatic carbocyclic ring system substituted with 0-4 R¹⁰ or R^{10a};

R³² is selected from:

- C(=O)-;
- C(=S)-
- S(=O)₂-;
- S(=O)-;
- P(=Z)(ZR¹³)-;

Z is S or O;

n'' and n' are independently 0-2;

R¹ and R²² are independently selected from the following groups:

- hydrogen,
- C₁-C₈ alkyl substituted with 0-2 R¹¹;
- C₂-C₈ alkenyl substituted with 0-2 R¹¹;

C₂-C₈ alkynyl substituted with 0-2 R¹¹;
C₃-C₁₀ cycloalkyl substituted with 0-2 R¹¹;

aryl substituted with 0-2 R¹²;

a 5-10-membered heterocyclic ring system containing
1-4 heteroatoms independently selected from N, S,
and O, said heterocyclic ring being substituted with
0-2 R¹²;

=O, F, Cl, Br, I, -CF₃, -CN, -CO₂R¹³, -C(=O)R¹³,
-C(=O)N(R¹³)₂, -CHO, -CH₂OR¹³, -OC(=O)R¹³,
-OC(=O)OR^{13a}, -OR¹³, -OC(=O)N(R¹³)₂, -NR¹³C(=O)R¹³,
-NR¹⁴C(=O)OR^{13a}, -NR¹³C(=O)N(R¹³)₂, -NR¹⁴SO₂N(R¹³)₂,
-NR¹⁴SO₂R^{13a}, -SO₃H, -SO₂R^{13a}, -SR¹³, -S(=O)R^{13a},
-SO₂N(R¹³)₂, -N(R¹³)₂, -NHC(=NH)NHR¹³, -C(=NH)NHR¹³,
=NOR¹³, NO₂, -C(=O)NHOR¹³, -C(=O)NHN(R¹³)R^{13a},
-OCH₂CO₂H, 2-(1-morpholino)ethoxy;

R¹ and R²¹ can alternatively join to form a 3-7 membered
carbocyclic ring substituted with 0-2 R¹²;

when n' is 2, R¹ or R²¹ can alternatively be taken
together with R¹ or R²¹ on an adjacent carbon atom to
form a direct bond, thereby to form a double or triple
bond between said carbon atoms;

R²² and R²³ can alternatively join to form a 3-7 membered
carbocyclic ring substituted with 0-2 R¹²;

when n" is 2, R²² or R²³ can alternatively be taken
together with R²² or R²³ on an adjacent carbon atom to
form a direct bond, thereby to form a double or triple
bond between the adjacent carbon atoms;

R¹ and R², where R²¹ is H, can alternatively join to form
a 5-8 membered carbocyclic ring substituted with 0-2 R¹²;

R¹¹ is selected from one or more of the following:

=O, F, Cl, Br, I, -CF₃, -CN, -CO₂R¹³, -C(=O)R¹³,
-C(=O)N(R¹³)₂, -CHO, -CH₂OR¹³, -OC(=O)R¹³,
-OC(=O)OR^{13a}, -OR¹³, -OC(=O)N(R¹³)₂, -NR¹³C(=O)R¹³,
-NR¹⁴C(=O)OR^{13a}, -NR¹³C(=O)N(R¹³)₂, -NR¹⁴SO₂N(R¹³)₂,
-NR¹⁴SO₂R^{13a}, -SO₃H, -SO₂R^{13a}, -SR¹³, -S(=O)R^{13a},
-SO₂N(R¹³)₂, -N(R¹³)₂, -NHC(=NH)NHR¹³, -C(=NH)NHR¹³,
=NOR¹³, NO₂, -C(=O)NHOR¹³, -C(=O)NHN(R¹³)R^{13a},
-OCH₂CO₂H, 2-(1-morpholino)ethoxy,

C₁-C₅ alkyl, C₂-C₄ alkenyl, C₃-C₆ cycloalkyl, C₃-C₆
cycloalkylmethyl, C₂-C₆ alkoxyalkyl, C₃-C₆
cycloalkoxy, C₁-C₄ alkyl (alkyl being substituted
with 1-5 groups selected independently from:
-NR¹³R¹⁴, -CF₃, NO₂, -SO₂R^{13a}, or -S(=O)R^{13a}),

aryl substituted with 0-2 R¹²,

a 5-10-membered heterocyclic ring system containing
1-4 heteroatoms independently selected from N, S,
and O, said heterocyclic ring being substituted with
0-2 R¹²;

R¹² is selected from one or more of the following:

phenyl, benzyl, phenethyl, phenoxy, benzyloxy,
halogen, hydroxy, nitro, cyano, C₁-C₅ alkyl, C₃-C₆
cycloalkyl, C₃-C₆ cycloalkylmethyl, C₇-C₁₀
arylalkyl, C₁-C₅ alkoxy, -CO₂R¹³, -C(=O)NHOR^{13a},
-C(=O)NHN(R¹³)₂, =NOR¹³, -B(R³⁴)(R³⁵), C₃-C₆
cycloalkoxy, -OC(=O)R¹³, -C(=O)R¹³, -OC(=O)OR^{13a},
-OR¹³, -(C₁-C₄ alkyl)-OR¹³, -N(R¹³)₂,
-OC(=O)N(R¹³)₂, -NR¹³C(=O)R¹³, -NR¹³C(=O)OR^{13a},
-NR¹³C(=O)N(R¹³)₂, -NR¹³SO₂N(R¹³)₂, -NR¹³SO₂R^{13a},
-SO₃H, -SO₂R^{13a}, -S(=O)R^{13a}, -SR¹³, -SO₂N(R¹³)₂,
C₂-C₆ alkoxyalkyl, methylenedioxy, ethylenedioxy,

C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, C₁-C₄ alkylcarbonyloxy, C₁-C₄ alkylcarbonyl, C₁-C₄ alkylcarbonylamino, -OCH₂CO₂H, 2-(1-morpholino)ethoxy, C₁-C₄ alkyl (alkyl being substituted with -N(R¹³)₂, -CF₃, NO₂, or -S(=O)R^{13a});

R¹³ is selected independently from: H, C₁-C₁₀ alkyl, C₃-C₁₀ cycloalkyl, C₄-C₁₂ alkylcycloalkyl, aryl, -(C₁-C₁₀ alkyl)aryl, or C₃-C₁₀ alkoxyalkyl;

R^{13a} is C₁-C₁₀ alkyl, C₃-C₁₀ cycloalkyl, C₄-C₁₂ alkylcycloalkyl, aryl, -(C₁-C₁₀ alkyl)aryl, or C₃-C₁₀ alkoxyalkyl;

when two R¹³ groups are bonded to a single N, said R¹³ groups may alternatively be taken together to form -(CH₂)₂₋₅- or -(CH₂)O(CH₂)-;

R¹⁴ is OH, H, C₁-C₄ alkyl, or benzyl;

R²¹ and R²³ are independently selected from:

hydrogen;

C₁-C₄ alkyl, optionally substituted with 1-6

halogen;

benzyl;

R² is H or C₁-C₈ alkyl;

R¹⁰ and R^{10a} are selected independently from one or more of the following:

phenyl, benzyl, phenethyl, phenoxy, benzyloxy, halogen, hydroxy, nitro, cyano, C₁-C₅ alkyl, C₃-C₆ cycloalkyl, C₃-C₆ cycloalkylmethyl, C₇-C₁₀ arylalkyl, C₁-C₅ alkoxy, -CO₂R¹³, -C(=O)N(R¹³)₂,

$-C(=O)NHOR^{13a}$, $-C(=O)NHN(R^{13})_2$, $=NOR^{13}$,
 $-B(R^{34})(R^{35})$, C_3-C_6 cycloalkoxy, $-OC(=O)R^{13}$,
 $-C(=O)R^{13}$, $-OC(=O)OR^{13a}$, $-OR^{13}$, $-(C_1-C_4 \text{ alkyl})-OR^{13}$,
 $-N(R^{13})_2$, $-OC(=O)N(R^{13})_2$, $-NR^{13}C(=O)R^{13}$,
 $-NR^{13}C(=O)OR^{13a}$, $-NR^{13}C(=O)N(R^{13})_2$, $-NR^{13}SO_2N(R^{13})_2$,
 $-NR^{13}SO_2R^{13a}$, $-SO_3H$, $-SO_2R^{13a}$, $-S(=O)R^{13a}$, $-SR^{13}$,
 $-SO_2N(R^{13})_2$, C_2-C_6 alkoxyalkyl, methylenedioxy,
 ethylenedioxy, C_1-C_4 haloalkyl (including $-C_vF_w$
 where $v = 1$ to 3 and $w = 1$ to $(2v+1)$), C_1-C_4
 haloalkoxy, C_1-C_4 alkylcarbonyloxy, C_1-C_4
 alkylcarbonyl, C_1-C_4 alkylcarbonylamino, $-OCH_2CO_2H$,
 $2-(1\text{-morpholino})ethoxy$, C_1-C_4 alkyl (alkyl being
 substituted with $-N(R^{13})_2$, $-CF_3$, NO_2 , or
 $-S(=O)R^{13a}$);

J is 3-aminopropionic acid or an L-isomer or D-isomer
 amino acid of structure $-N(R^3)C(R^4)(R^5)C(=O)-$, wherein:

R^3 is H or C_1-C_8 alkyl;

R^4 is H or C_1-C_3 alkyl;

R^5 is selected from:

hydrogen;

C_1-C_8 alkyl substituted with 0-2 R^{11} ;

C_2-C_8 alkenyl substituted with 0-2 R^{11} ;

C_2-C_8 alkynyl substituted with 0-2 R^{11} ;

C_3-C_{10} cycloalkyl substituted with 0-2 R^{11} ;


aryl substituted with 0-2 R^{12} ;

a 5-10-membered heterocyclic ring system containing
 1-4 heteroatoms independently selected from N, S, or
 O, said heterocyclic ring being substituted with 0-2
 R^{12} ;

$=O$, F, Cl, Br, I, $-CF_3$, $-CN$, $-CO_2R^{13}$, $-C(=O)R^{13}$,


$-\text{C}(=\text{O})\text{N}(\text{R}^{13})_2$, $-\text{CHO}$, $-\text{CH}_2\text{OR}^{13}$, $-\text{OC}(=\text{O})\text{R}^{13}$,
 $-\text{OC}(=\text{O})\text{OR}^{13a}$, $-\text{OR}^{13}$, $-\text{OC}(=\text{O})\text{N}(\text{R}^{13})_2$, $-\text{NR}^{13}\text{C}(=\text{O})\text{R}^{13}$,
 $-\text{NR}^{14}\text{C}(=\text{O})\text{OR}^{13a}$, $-\text{NR}^{13}\text{C}(=\text{O})\text{N}(\text{R}^{13})_2$, $-\text{NR}^{14}\text{SO}_2\text{N}(\text{R}^{13})_2$,
 $-\text{NR}^{14}\text{SO}_2\text{R}^{13a}$, $-\text{SO}_3\text{H}$, $-\text{SO}_2\text{R}^{13a}$, $-\text{SR}^{13}$, $-\text{S}(=\text{O})\text{R}^{13a}$,
 $-\text{SO}_2\text{N}(\text{R}^{13})_2$, $-\text{N}(\text{R}^{13})_2$, $-\text{NHC}(=\text{NH})\text{NHR}^{13}$, $-\text{C}(=\text{NH})\text{NHR}^{13}$,
 $=\text{NOR}^{13}$, NO_2 , $-\text{C}(=\text{O})\text{NHOR}^{13}$, $-\text{C}(=\text{O})\text{NHNHR}^{13}\text{R}^{13a}$, $=\text{NOR}^{13}$,
 $-\text{B}(\text{R}^{34})(\text{R}^{35})$, $-\text{OCH}_2\text{CO}_2\text{H}$, 2-(1-morpholino)ethoxy,
 $-\text{SC}(=\text{NH})\text{NHR}^{13}$, N_3 , $-\text{Si}(\text{CH}_3)_3$, $(\text{C}_1\text{-C}_5 \text{ alkyl})\text{NHR}^{16}$;

$-(\text{C}_0\text{-C}_6 \text{ alkyl})\text{X}$;



, where q is

independently 0,1;





;

$-(\text{CH}_2)_m\text{S}(\text{O})_{p'}(\text{CH}_2)_2\text{X}$, where $m = 1, 2$ and $p' = 0-2$;

wherein X is defined below; and

R^3 and R^4 may also be taken together to form





;

R³ and R⁵ can alternatively be taken together to form
 -(CH₂)_t- or -CH₂S(O)_pC(CH₃)₂-, where t = 2-4 and p' =
 0-2; or

R⁴ and R⁵ can alternatively be taken together to form
 -(CH₂)_u-, where u = 2-5;

R¹⁶ is selected from:


- an amine protecting group;
- 1-2 amino acids;
- 1-2 amino acids substituted with an amine protecting group;

K is a D-isomer or L-isomer amino acid of structure
 -(R⁶)CH(R⁷)C(=O)-, wherein:

R⁶ is H or C₁-C₈ alkyl;

R⁷ is selected from:

-(C₁-C₇ alkyl)X;




, wherein each q is

independently 0-2 and substitution on the phenyl is at the 3 or 4 position;



, wherein each


q is independently 0-2 and substitution on the cyclohexyl is at the 3 or 4 position;



$-(CH_2)_mO-(C_1-C_4 \text{ alkyl})-X$, where $m = 1$ or 2 ;

$-(CH_2)_mS(O)_{p'}-(C_1-C_4 \text{ alkyl})-X$, where $m = 1$ or 2 and $p' = 0-2$; and


X is selected from:




;

$-N(R^{13})R^{13}$; $-C(=NH)(NH_2)$; $-SC(=NH)-NH_2$; $-NH-C(=NH)(NHCN)$;
 $-NH-C(=NCN)(NH_2)$; $-NH-C(=N-OR^{13})(NH_2)$;

R⁶ and R⁷ can alternatively be taken together to form



or 2 and wherein n = 0 or 1 and X is -NH₂ or




;

L is -Y(CH₂)_vC(=O)-, wherein:

Y is NH, N(C₁-C₃ alkyl), O, or S; and v = 1 or 2;

M is a D-isomer or L-isomer amino acid of structure



, wherein:

q' is 0-2;

R¹⁷ is H, C₁-C₃ alkyl;

R⁸ is selected from:

-CO₂R¹³, -SO₃R¹³, -SO₂NHR¹⁴, -B(R³⁴)(R³⁵), -NHSO₂CF₃,
-CONHNHSO₂CF₃, -PO(OR¹³)₂, -PO(OR¹³)R¹³,
-SO₂NH-heteroaryl (said heteroaryl being
5-10-membered and having 1-4 heteroatoms selected
independently from N, S, or O), -SO₂NH-heteroaryl
(said heteroaryl being 5-10-membered and having 1-4
heteroatoms selected independently from N, S, or O),
-SO₂NHCOR¹³, -CONHSO₂R^{13a}, -CH₂CONHSO₂R^{13a},
-NHSO₂NHCOR^{13a}, -NHCONHSO₂R^{13a}, -SO₂NHCONHR¹³;

R³⁴ and R³⁵ are independently selected from:

-OH,
-F,
-N(R¹³)₂, or
C₁-C₈-alkoxy;


R³⁴ and R³⁵ can alternatively be taken together form:

a cyclic boron ester where said chain or ring
contains from 2 to 20 carbon atoms and, optionally,
1-4 heteroatoms independently selected from N, S, or
O;

a divalent cyclic boron amide where said chain or
ring contains from 2 to 20 carbon atoms and,
optionally, 1-4 heteroatoms independently selected
from N, S, or O;

a cyclic boron amide-ester where said chain or ring
contains from 2 to 20 carbon atoms and, optionally,
1-4 heteroatoms independently selected from N, S, or
O.

7. The method of Claim 6 wherein the localization step
comprises the step of localizing a compound of the
formula (I) at the thrombus wherein Q is of the formula
(III),



(III)

or a pharmaceutically acceptable salt or prodrug form thereof wherein:

the shown phenyl ring may be further substituted with 0-3 R¹⁰;

R¹⁰ is selected independently from: H, C₁-C₈ alkyl, phenyl, halogen, or C₁-C₄ alkoxy;

R¹ is H, C₁-C₄ alkyl, phenyl, benzyl, or phenyl-(C₁-C₄)alkyl;

R² is H or methyl;

R¹³ is selected independently from: H, C₁-C₁₀ alkyl, C₃-C₁₀ cycloalkyl, C₄-C₁₂ alkylcycloalkyl, aryl, -(C₁-C₁₀ alkyl)aryl, or C₃-C₁₀ alkoxyalkyl;

R^{13a} is C₁-C₁₀ alkyl, C₃-C₁₀ cycloalkyl, C₄-C₁₂ alkylcycloalkyl, aryl, -(C₁-C₁₀ alkyl)aryl, or C₃-C₁₀ alkoxyalkyl;

when two R^{13} groups are bonded to a single N, said R^{13} groups may alternatively be taken together to form $-(CH_2)_2-$ or $-(CH_2)O(CH_2)-$;

R^{14} is OH, H, C_1 - C_4 alkyl, or benzyl;

J is β -alanine or an L-isomer or D-isomer amino acid of structure $-N(R^3)C(R^4)(R^5)C(=O)-$, wherein:

R^3 is H or CH_3 ;

R^4 is H or C_1 - C_3 alkyl;

R^5 is H, C_1 - C_8 alkyl, C_3 - C_6 cycloalkyl, C_3 - C_6 cycloalkylmethyl, C_1 - C_6 cycloalkylethyl, phenyl, phenylmethyl, CH_2OH , CH_2SH , CH_2OCH_3 , CH_2SCH_3 , $CH_2CH_2SCH_3$, $(CH_2)_sNH_2$, $-(CH_2)_sNHC(=NH)(NH_2)$, $-(CH_2)_sNHR^{16}$, where $s = 3-5$; or

R^{16} is selected from:

an amine protecting group;

1-2 amino acids; or

1-2 amino acids substituted with an amine protecting group;


R^3 and R^5 can alternatively be taken together to form $-CH_2CH_2CH_2-$; or

R^4 and R^5 can alternatively be taken together to form $-(CH_2)_u-$, where $u = 2-5$;




K is an L-isomer amino acid of structure $-N(R^6)CH(R^7)C(=O)-$, wherein:

R^6 is H or C_1 - C_8 alkyl;

R^7 is:





;




;



;






$-(\text{CH}_2)_m\text{O}-(\text{C}_1\text{---C}_4 \text{ alkyl})\text{---NH}-(\text{C}_1\text{---C}_6 \text{ alkyl})$, where $m = 1$ or 2 ;

$-(\text{CH}_2)_m\text{S}-(\text{C}_1\text{---C}_4 \text{ alkyl})\text{---NH}-(\text{C}_1\text{---C}_6 \text{ alkyl})$, where $m = 1$ or 2 ; and

X is $-\text{NH}_2$ or $-\text{NHC}(=\text{NH})(\text{NH}_2)$, provided that X is not $-\text{NH}_2$ when $r = 4$; or

R^6 and R^7 are alternatively be taken together to form




$-\text{NHC}(=\text{NH})(\text{NH}_2)$;

L is $-\text{Y}(\text{CH}_2)_v\text{C}(=\text{O})\text{---}$, wherein:

Y is NH, O, or S; and $v = 1, 2$;

M is a D-isomer or L-isomer amino acid of structure



, wherein:


q' is 0-2;

R¹⁷ is H, C₁-C₃ alkyl;

R⁸ is selected from:

-CO₂R¹³, -SO₃R¹³, -SO₂NHR¹⁴, -B(R³⁴)(R³⁵), -NHSO₂CF₃,
-CONHNHSO₂CF₃, -PO(OR¹³)₂, -PO(OR¹³)R¹³,
-SO₂NH-heteroaryl (said heteroaryl being
5-10-membered and having 1-4 heteroatoms selected
independently from N, S, or O), -SO₂NH-heteroaryl
(said heteroaryl being 5-10-membered and having 1-4
heteroatoms selected independently from N, S, or O),
-SO₂NHCOR¹³, -CONHSO₂R^{13a}, -CH₂CONHSO₂R^{13a},
-NHSO₂NHCOR^{13a}, -NHCONHSO₂R^{13a}, -SO₂NHCONHR¹³.


8. The method of Claim 4 wherein the localization step comprises the step of localizing a compound of the formula (IV) at the thrombus:



(IV).

9. The method of Claim 4 wherein the localization step

comprises the step of localizing a compound of the formula (V) at the thrombus:



(V).

10. The method of Claim 1 wherein the acquisition step comprises the step of acquiring image slices representing a concentration of radioactivity associated with the thrombus.
11. The method of Claim 10 wherein the acquisition step comprises the step of acquiring single photon emission computed tomography images of the thrombus.
12. The method of Claim 1 wherein the acquisition step comprises the step of acquiring transaxial image slices and further comprising the step of reformatting the transaxial image slices into image slices that are parallel to a long axis associated with the thrombus.
13. The method of Claim 1 comprising the step of displaying the two-dimensional array as a reprojected image.


14. The method of Claim 1 wherein the scanning step is performed at a series of angles.
15. The method of Claim 14 wherein the assignment step is performed at each of the series of angles.
16. The method of Claim 15 comprising the step of sequentially displaying the two-dimensional arrays as reprojected images.
17. A method for imaging a pulmonary embolus comprising the steps of:
 - a. localizing a radiolabelled compound at the pulmonary embolus;
 - b. acquiring image slices representing a physical property of the radiolabelled pulmonary embolus;
 - c. assembling the image slices into a three-dimensional matrix of data;
 - d. scanning the three-dimensional matrix of data along an array of parallel lines to determine a maximum value along each line; and
 - e. assigning the maximum value along each line to a pixel in a two-dimensional array, the position of the pixel corresponding to the position of the line in the array of parallel lines.
18. A method for imaging an arterial thrombus comprising the steps of:
 - a. localizing a radiolabelled compound at the arterial thrombus;
 - b. acquiring image slices representing a physical property of the radiolabelled arterial thrombus;
 - c. assembling the image slices into a three-dimensional matrix of data;
 - d. scanning the three-dimensional matrix of data along an array of parallel lines to determine a maximum

- value along each line; and
- e. assigning the maximum value along each line to a pixel in a two-dimensional array, the position of the pixel corresponding to the position of the line in the array of parallel lines.
19. A method for imaging a coronary thrombus comprising the steps of:
- a. localizing a radiolabelled compound at the coronary thrombus;
 - b. acquiring image slices representing a physical property of the radiolabelled coronary thrombus;
 - c. assembling the image slices into a three-dimensional matrix of data;
 - d. scanning the three-dimensional matrix of data along an array of parallel lines to determine a maximum value along each line; and
 - e. assigning the maximum value along each line to a pixel in a two-dimensional array, the position of the pixel corresponding to the position of the line in the array of parallel lines.

ABSTRACT OF THE DISCLOSURE

A method wherein a radiolabelled compound is localized at a thrombus. Two-dimensional images, representing a physical property associated with the radiolabelled thrombus, are acquired and assembled into a three-dimensional matrix of data. The three-dimensional matrix of data is then scanned along an array of parallel lines to determine a maximum value along each line. The maximum value along each line is then assigned to a pixel in a two-dimensional array, where the relative position of the pixel in the two-dimensional array corresponds to the relative position of the line in the array of parallel lines.

Fig. 1



DECLARATION and POWER OF ATTORNEY

As a below-named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

METHOD FOR LOCALIZATION OF BLOOD CLOTS

the specification of which is attached hereto under the following box is checked:

☐ was filed on _____ as U.S. Application No. _____ or PCT International Application No. _____ and was amended on _____ (if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is known to me to be material to patentability as defined in 37 CFR § 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed.

| Application No. | Country | Filing Date | Priority Claimed (Yes/No) |
|-----------------|---------|-------------|---------------------------|
|-----------------|---------|-------------|---------------------------|

I hereby claim the benefit under 35 U.S.C. § 119(e) of any United States Provisional Application(s) listed below.

U.S. Provisional Application No.
60/126,359

U.S. Filing Date
3/26/99

I hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s), or § 365(c) of any PCT International Application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application or PCT International Application in the manner provided by the first paragraph of 35 U.S.C. § 112, I acknowledge the duty to disclose information which is known to me to be material to patentability as defined in 37 CFR § 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application.

| Application No. | U.S. Filing Date | Status (patented, pending or abandoned) |
|-----------------|------------------|---|
|-----------------|------------------|---|

POWER OF ATTORNEY: I hereby appoint the following attorney(s) and/or agent(s) the power to prosecute this application and transact all business in the Patent and Trademark Office connected therewith:

| | |
|---|---|
| Name: Blair Q. Ferguson Gerald J. Boudreaux Karen H. Kondrad Scott K. Larsen Maureen P. O'Brien Norbert Reinert Mary K. VanAtten Kenneth B. Rubin Rosemarie R. Wilk-Orescan | Registration No.: 34,329 35,073 38,212 38,532 42,043 18,926 39,408 36,295 P45,220 |
|---|---|

| | | |
|--|--|--|
| Send correspondence and direct telephone calls to: Maureen P. O'Brien | DuPont Pharmaceuticals Company c/o E. I. du Pont de Nemours and Co. Legal - Patents 1007 Market Street Wilmington, DE 19898, U.S.A. | Tel. No. (302) 992-4528 |
|--|--|--|

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

| INVENTOR(S) | | | |
|------------------------------------|---------------------|--------------------------|------------------------|
| Full Name of Inventor | Last Name | First Name | Middle Name |
| | LAZEWATSKY | JOEL | |
| Signature (please sign full name): | | | Date: |
| Residence & Citizenship | City | State or Foreign Country | Country of Citizenship |
| | AUBURDALE | MASSACHUSETTS | US |
| Post Office Address | Post Office Address | City | State or Country |
| | 32 WOODLAND RD. | AUBURDALE | MASSACHUSETTS |
| | | | Zip Code |
| | | | 02466 |
| Full Name of Inventor | Last Name | First Name | Middle Name |
| Signature (please sign full name): | | | Date: |
| Residence & Citizenship | City | State or Foreign Country | Country of Citizenship |
| | | | |
| Post Office Address | Post Office Address | City | State or Country |
| | | | Zip Code |
| | | | |

☐ Additional Inventors are being named on separately numbered sheets attached hereto.